

Contents

Foreword	vii
Acknowledgements	ix
1 Introduction: investigating the archaeology of the southern Curtis Coast	1
Introduction	1
The continental narrative	1
Refining mid-to-late Holocene chronologies	5
Rockshelters versus open sites: some sampling issues	7
Rockshelters versus open sites: some taphonomic issues	8
Background to the study	10
The study region	10
Aims of the study	11
2 The study region: the southern Curtis Coast	13
Introduction	13
Physical setting	13
Geology and geomorphology	14
Climate	17
Hydrology	17
Flora	18
Fauna	18
Palaeoenvironment and environmental change	20
Cultural setting	22
Previous archaeological research	25
Gooreng Gooreng Cultural Heritage Project	27
Other dated archaeological sites in the region	30
Agnes Beach Midden (ABM)	30
Middle Island Sandblow Site (MISS)	31
Round Hill Creek Mound (RHCM)	33
Worthington Creek Midden (WCM)	33
Discussion	34
Summary	35
3 Methods of investigation	37
Introduction	37

Excavation strategy	37
Excavation methods	38
Radiocarbon dating and calibration	39
Marine and estuarine reservoir effects	39
Radiocarbon age calibration	40
Dating terminology	40
Laboratory analyses	41
Invertebrate remains	41
Vertebrate remains	44
Stone artefacts	45
Non-artefactual stone	45
Charcoal	45
Organic material	45
Age-depth curves and analytical units	45
Summary	45
4 Marine and estuarine reservoir effects in central Queensland: determination of ΔR values	47
Introduction	47
Background	49
Australian marine reservoir studies: a review	51
The original study	51
Further studies	53
The present study: methods	55
Results	56
Live-collected known-age samples	56
Archaeological shell/charcoal paired samples	57
Temporal variability or $\Delta R(t)$	61
Discussion	61
Summary	64
5 Bivalve conjoin analyses: assessing site integrity	65
Introduction	65
Background	65
General methods and approach	68
Seven Mile Creek Mound bivalve conjoin analysis	69
Introduction	69
Aims	69
Methods	70
Results	70
Bivalve conjoin analysis blind test	71
Introduction	71
Aims	71
Methods	71
Results	72
Discussion	72
Summary	77
6 Seven Mile Creek Mound	79
Introduction	79

	Site description and setting	79
	Excavation methods	81
	Cultural deposit and stratigraphy	82
	Radiocarbon dating and chronology	83
	Stratigraphic integrity and disturbance	84
	Laboratory methods	85
	Cultural materials	85
	Invertebrate remains	85
	Vertebrate remains	92
	Stone artefacts	92
	Other remains	93
	Discussion	94
	Summary	96
7	Mort Creek Site Complex	97
	Introduction	97
	Site description and setting	97
	Previous investigations	99
	Excavation methods	101
	Cultural deposit and stratigraphy	102
	Radiocarbon dating and chronology	102
	Stratigraphic integrity and disturbance	104
	Laboratory methods	106
	Cultural materials	106
	Invertebrate remains	106
	Vertebrate remains	110
	Stone artefacts	111
	Other remains	112
	Discussion	112
	Summary	113
8	Pancake Creek Site Complex	115
	Introduction	115
	Site description and setting	115
	Excavation methods	117
	Cultural deposit and stratigraphy	119
	Radiocarbon dating and chronology	122
	Stratigraphic integrity and disturbance	123
	Laboratory methods	123
	Cultural materials	124
	Invertebrate remains	124
	Vertebrate remains	124
	Stone artefacts	124
	Other remains	128
	Discussion	129
	Summary	129
9	Ironbark Site Complex	131
	Introduction	131

Site description and setting	131
Excavation methods	135
Cultural deposit and stratigraphy	139
Radiocarbon dating and chronology	144
Stratigraphic integrity and disturbance	146
Laboratory methods	147
Cultural materials	147
Invertebrate remains	147
Vertebrate remains	151
Stone artefacts	152
Glass artefacts	152
Other remains	154
Discussion	154
Summary	156
10 Eurimbula Creek 1	157
Introduction	157
Site description and setting	157
Excavation methods	160
Cultural deposit and stratigraphy	160
Radiocarbon dating and chronology	161
Stratigraphic integrity and disturbance	162
Laboratory methods	162
Cultural materials	162
Invertebrate remains	162
Vertebrate remains	163
Other remains	166
Discussion	166
Summary	167
11 Eurimbula Creek 2	169
Introduction	169
Site description and setting	169
Excavation methods	171
Cultural deposit and stratigraphy	171
Radiocarbon dating and chronology	172
Stratigraphic integrity and disturbance	173
Laboratory methods	173
Cultural materials	173
Invertebrate remains	173
Other remains	174
Discussion	176
Summary	176
12 Eurimbula Site 1	177
Introduction	177
Site description and setting	177
Previous investigations	180
Excavation methods	184

	Cultural deposit and stratigraphy	184
	Radiocarbon dating and chronology	186
	Stratigraphic integrity and disturbance	188
	Laboratory methods	189
	Cultural materials	189
	Invertebrate remains	189
	Vertebrate remains	194
	Stone artefacts	196
	Other remains	197
	Discussion	197
	Summary	199
13	Tom's Creek Site Complex	201
	Introduction	201
	Site description and setting	201
	Excavation methods	204
	Cultural deposit and stratigraphy	207
	Radiocarbon dating and chronology	209
	Stratigraphic integrity and disturbance	210
	Laboratory methods	212
	Cultural materials	212
	Invertebrate remains	212
	Vertebrate remains	218
	Stone artefacts	220
	Glass artefacts	222
	Other remains	223
	Discussion	223
	Summary	224
14	Synthesis of results: towards an archaeology of the southern Curtis Coast	225
	Introduction	225
	Regional site chronology	226
	Regional site contents	232
	Shellfish remains	232
	Fish remains	234
	Charcoal	236
	Stone artefacts	236
	Regional patterns in site structure	238
	Discussion	240
	Modelling regional settlement histories	241
	Phase I (pre-4,000 BP–1,500 BP)	241
	Phase II (c.1,500 BP–c.AD 1850s)	243
	Phase III (c.AD 1850s–c.AD 1920s)	244
	Summary	245
15	Wider implications and conclusions	247
	Introduction	247
	Key findings	247
	Regional context and implications	248

Regional occupation before the late Holocene	248
Reduced occupation or abandonment of the coastal zone in the late Holocene	250
The antiquity of marine fishing	251
Localisation of resource use	251
Wider implications	252
Methodological implications	254
Dating strategies and chronological control	254
Integrity of open sites	254
Discussion	255
Conclusion	255
References	257
Appendices	279
Appendix 1	280
Radiocarbon dates: technical data	
Appendix 2	283
Recorded archaeological sites on the southern Curtis Coast	
Appendix 3	293
Site name synonyms for recorded sites on the southern Curtis Coast	
Appendix 4	297
Excavation data	
Appendix 5	313
Shellfish reference collection	

List of Figures

1.1	The Australian region, showing places mentioned in the text (after Allen and O'Connell 1995:vi).	5
1.2	Distribution of dated archaeological sites in Queensland with a Holocene component (Ulm and Reid 2000:3).	8
2.1	The Gooreng Gooreng Cultural Heritage Project study area, showing major towns and the general distribution of Gooreng Gooreng speakers (heavy line) (after Horton 1994; Williams 1981). The southern Curtis Coast study area is shown by the box.	15
2.2	The southern Curtis Coast study area, showing all recorded archaeological sites as triangles (after Ulm and Lilley 1999). Site designations are shown for sites which are not illustrated in Figs 2.9–2.11. Heavy black lines on Middle Island indicate the general location of extensive low density shell deposits.	16
2.3	General view of estuary and near-coastal ranges across Jenny Lind Creek. Such estuaries are common in the region and support diverse populations of marine life. Facing southwest.	19
2.4	Freshwater wetlands adjoining the upper reaches of Round Hill Creek. This swamp is part of a vast network of interconnected wetland areas located in the swales between ridge systems. Facing north.	19
2.5	Example of recent bank recession of the north bank of Middle Creek, showing concrete stairs once used to access the beach from the top of the erosion bank some 15m away from the contemporary section. Facing northwest.	19
2.6	Beach ridge vegetation in Eurimbula National Park, including weeping cabbage palm communities (<i>L. australis</i>) and bracken (<i>P. esculentum</i>).	20
2.7	Mud ark (<i>A. trapezia</i>) in a dense paddock of seagrass (<i>Z. capricornia</i>) in the middle of Round Hill Creek. The presence of mud ark valves in sites dated to 4,000 BP indicate that seagrass beds are a long-term feature of the local environment. Facing west.	20
2.8	Cluster of telescope mud whelk (<i>T. telescopium</i>) on the mangrove fringe of Eurimbula Creek. These whelks are abundant throughout the region, although virtually absent from archaeological deposits. Facing west.	20
2.9	Northern segment of the southern Curtis Coast study area showing recorded archaeological sites as triangles in the Hummock Hill Island and Seven Mile Creek areas. EPA site numbers are shown. Sites without a 'JE:' prefix have a 'KE:' prefix which has been omitted owing to space limitations.	28
2.10	Central segment of the southern Curtis Coast study area showing recorded archaeological sites as triangles in the Turkey Beach, Rodds Peninsula and Middle Island areas. EPA site numbers are shown where available. Sites without a 'JF:' prefix have a 'KE:' prefix which has been omitted owing to space limitations. Sites with a 'SCC' prefix are currently unregistered sites.	28
2.11	Southern segment of the southern Curtis Coast study area showing recorded archaeological sites as triangles in the Agnes Water, Round Hill Creek, Eurimbula Creek and Middle Creek areas. EPA site numbers are shown. All sites have a 'KE:' prefix which has been omitted owing to space limitations.	29
2.12	Stone-walled tidal fishtrap at Richards Point on Rodds Peninsula. This is one of two such features recorded in the study region. The second fishtrap is located in Mort Creek 4km to the south (see Chapter 8). Facing north (Photograph: Ian Lilley).	31

2.13	Scarred tree on eucalypt at Agnes Water (KE:A60) (Photograph: Environmental Protection Agency).	31
2.14	Agnes Beach Midden. Facing southwest.	32
2.15	Worthington Creek Midden. Facing northwest.	32
2.16	Middle Island Sandblow Site, showing pipi (<i>D. deltoides</i>) scatter on the surface of the sandblow. A range of cultural and non-cultural material is exposed on the eastern margin of the sandblow. Facing north.	32
2.17	Middle Island Sandblow Site, showing microgranite grinding implement located on the surface of the sandblow. Obviously modified and apparently unmodified pieces of stone are common along the eastern margin of the sandblow. Facing south.	32
3.1	Bivalve terminology (after Claassen 1998:21; Hedley 1904).	42
3.2	Bivalve terminology and measured attributes (after Claassen 1998:21; Hedley 1904).	42
3.3	Gastropod terminology (after Andrews 1971:42; Claassen 1998:19).	43
4.1	Map of Australia, showing places mentioned in the text. The boxed area indicates the southern Curtis Coast study area.	48
4.2	(a) Part of the central Queensland coast showing the location of 'pre-bomb' live-collected shell specimens (☒) and (b) estuaries and archaeological sites (▲) discussed in the text. SMCM=Seven Mile Creek Mound; MCSC=Mort Creek Site Complex; PCSC=Pancake Creek Site Complex; ISC=Ironbark Site Complex; ES1=Eurimbula Site 1; TCSC=Tom's Creek Site Complex; ABM=Agnes Beach Midden.	49
4.3	Example of ΔR calculation method for pair NZA-12117/Wk-8326 (see caption for Table 4.6).	60
4.4	ΔR values calculated for the southern Curtis Coast.	63
4.5	Calibrated radiocarbon age-ranges from the Seven Mile Creek Mound, using various ΔR values (see Table 4.7).	63
5.1	Schematic representation of conjoin identification procedure. O=individual <i>A. trapezia</i> valve. In this matrix, left and right valves are arranged in descending size order. See text for details.	71
5.2	Maximum vertical distance separating all conjoined valve-pairs, Seven Mile Creek Mound, Square A.	75
5.3	Distribution of identified <i>A. trapezia</i> valve-pairs (n=61), Seven Mile Creek Mound, Square A. An additional 22 valve-pairs encountered as articulated specimens during excavation are shown as short horizontal lines down the right hand side of the figure (see Chapter 6). Line termination points indicate the vertical mid-points of the excavation units from which conjoining valves were recovered. Short horizontal lines indicate valve-pairs identified within excavation units. Not to scale on the horizontal axis.	75
5.4	Particle size distribution of <i>A. trapezia</i> after heating at various temperatures and then mechanical destruction for 30 seconds (after Robins and Stock 1990:98).	76
5.5	Fragmentation of <i>A. trapezia</i> , Seven Mile Creek Mound, Square A, expressed as the number of fragments per 100g of shell.	76
6.1	The Seven Mile Creek catchment area showing the location of the Seven Mile Creek Mound as a triangle. Dark grey shading indicates the general extent of mangrove, saltflats and claypans.	80
6.2	Topographic map of the site area, showing the position of the excavation. The site datum is shown as a triangle. Dashed lines indicate the location of the tide strand line during the period of excavation. The wide shaded line denotes a 4WD track on the edge of the saltflats. Contours in 10cm intervals.	80
6.3	General view of the Seven Mile Creek Mound. Note the low fringing mangroves on the intertidal flats through the gap in trees at rear left of frame. Facing south.	81

6.4	General view of the completed excavation. Note the dark zone of grass root penetration at the top of the deposit. Facing north.	81
6.5	General view of completed excavation (Squares D–A). Note the stratigraphic break at the base of the shell deposit. Facing north.	81
6.6	Stratigraphic section, Seven Mile Creek Mound, Squares A–D.	82
6.7	Distribution of identified <i>A. trapezia</i> valve-pairs (n=61), Seven Mile Creek Mound, Square A. An additional 22 valve-pairs encountered as articulated specimens during excavation are shown as short horizontal lines down the right hand side of the figure. Line termination points indicate the vertical mid-points of the excavation units from which conjoining valves were recovered. Short horizontal lines indicate valve-pairs identified within excavation units. Not to scale on the horizontal axis. See Chapter 5 for details of methods.	85
6.8	Abundance of oyster (<i>S. glomerata</i>).	86
6.9	Abundance of hairy mussel (<i>T. hirsutus</i>).	86
6.10	Abundance of mud ark (<i>A. trapezia</i>).	87
6.11	Abundance of scallop (<i>P. sugillata</i>).	87
6.12	Abundance of fish bone.	87
6.13	Abundance of charcoal.	90
6.14	Abundance of artefactual stone.	90
6.15	Abundance of non-artefactual stone.	90
6.16	Relative contribution of dominant shellfish taxa. Note that XU1 is excluded as it contained negligible quantities of shell (see Appendix 4).	91
7.1	The Mort Creek catchment area. Dark grey shading indicates the general extent of mangrove, saltflats and claypans. Dashed lines denote 4WD tracks.	99
7.2	Topographic map of the Mort Creek Site Complex. Contours are in 0.5m intervals. Dark grey shading indicates the general extent of mangroves. Dots indicate auger test holes (not all shown).	99
7.3	Chenier ridge at the Mort Creek Site Complex extending into Rodds Harbour. Facing southwest (Photograph: Ian Lilley).	101
7.4	General view of completed excavation, Squares A–D. Note continuous microgranite bedrock across the base of the excavation. Facing north.	101
7.5	General view of completed excavation, showing the section of Squares C–D. Note shell layer across the upper 20cm of the deposit. Facing east.	101
7.6	Stratigraphic section, Mort Creek Site Complex, Squares A–D.	103
7.7	Distribution of identified <i>A. trapezia</i> valve-pairs (n=10), Mort Creek Site Complex, Square C. Line termination points indicate the vertical mid-points of the excavation units from which conjoining valves were located. Short horizontal lines indicate valve-pairs identified within excavation units. Not to scale on the horizontal axis. See Chapter 5 for details of methods.	106
7.8	Abundance of mud ark (<i>A. trapezia</i>).	107
7.9	Abundance of oyster (<i>S. glomerata</i>).	107
7.10	Abundance of whelk (<i>P. ebininus</i>).	108
7.11	Abundance of fish bone.	108
7.12	Abundance of organic material.	108
7.13	Abundance of charcoal.	109
7.14	Abundance of artefactual stone.	109
7.15	Abundance of non-artefactual stone.	109
8.1	The Pancake Creek catchment area showing the location of the Pancake Creek Site Complex. Dark grey shading indicates the general extent of mangrove, saltflats and claypans.	117

8.2	Detail of the site area showing the location of excavated squares and beach ridge vegetation units. 3a= <i>Corymbia intermedia</i> tall woodland with shrub/heath midstratum; 3b= <i>C. tessellaris</i> , <i>C. intermedia</i> tall to very tall woodland with <i>Livistona decipiens</i> / <i>Melaleuca dealbata</i> and shrub understorey; 3c= <i>L. decipiens</i> / <i>Melaleuca</i> tall forest/tall open forest with emergent <i>Eucalyptus tereticornis</i> / <i>C. intermedia</i> ; 3d= <i>L. decipiens</i> open forest; 3e= <i>Acacia julifera</i> / <i>A. flavescens</i> tall shrubland; 3f= <i>M. dealbata</i> open forest; 5a= <i>C. tessellaris</i> woodland with understorey of low microphyll vine thicket; 5c= <i>Casuarina equisetifolia</i> low open woodland; 5d=Tall open shrubland (windsheared vegetation) (after QDEH 1997).	117
8.3	Eroding bank in the vicinity of Squares G–H showing displaced trees and shell. Facing southwest.	118
8.4	General view of completed excavation at Squares C–F. Facing south.	119
8.5	Close-up view of concentration of <i>A. trapezia</i> valves encountered in Square H, XU9, at a depth of 26–30cm. These shells date to around 600 years ago and include a single valve conjoin. Facing southeast.	119
8.6	Stratigraphic section, Pancake Creek Site Complex, Squares A–B.	120
8.7	Stratigraphic section, Pancake Creek Site Complex, Squares C–F.	121
8.8	Stratigraphic section, Pancake Creek Site Complex, Squares G–H.	121
8.9	Abundance of oyster (<i>S. glomerata</i>).	126
8.10	Abundance of mud ark (<i>A. trapezia</i>).	126
8.11	Abundance of whelk (<i>P. ebininus</i>).	126
8.12	Abundance of scallop (<i>P. sugillata</i>).	127
8.13	Abundance of hairy mussel (<i>T. hirsutus</i>).	127
8.14	Abundance of organic material.	127
8.15	Abundance of charcoal.	128
8.16	Abundance of non-artefactual stone.	128
9.1	The Middle Creek catchment area showing the location of the Ironbark Site Complex (ISC) and nearby excavated sites (EC1=Eurimbula Creek 1; EC2=Eurimbula Creek 2; ES1=Eurimbula Site 1; TCSC=Tom’s Creek Site Complex). Dark grey shading indicates the general extent of mangrove, saltflats and claypans. Dotted shading indicates land above 200m. Solid dots indicate local population centres.	133
9.2	Aerial view of the Ironbark Site Complex, showing the maximum extent from the creek of shell and stone exposed at the surface (heavy line) and the general location of excavation squares (after BPA Run 15D/79, 30 July 1996). Based on data provided by the Department of Natural Resources and Mines, Queensland 2006, which gives no warranty in relation to the data (including accuracy, reliability, completeness or suitability) and accepts no liability (including without limitation, liability in negligence) for any loss, damage or costs (including consequential damage) relating to any use of the data.	134
9.3	Site plan of Ironbark Site Complex, showing area of Squares A–M. Contours are in 0.5m intervals. Rock outcrops and low density artefactual material continue for a further c.50m southwest of the limits of this plan.	135
9.4	Site plan of Ironbark Site Complex, showing area of Square N. Contours are in 0.5m intervals. Only cycads over 50cm in trunk height are shown.	136
9.5	Site plan of Ironbark Site Complex, showing area of Squares O–P. Contours are in 0.5m intervals. Tree stumps are shown on the fringing beach.	137
9.6	Site plan of Ironbark Site Complex, showing area of Squares Q–R. Contours are in 0.5m intervals.	137
9.7	General view of the western side of the quarry sloping into Middle Creek, showing a massive core in centre foreground. Middle Island in background. Facing north.	139
9.8	Surface of Square E, XU1, showing photographic recording method. Facing north.	139

9.9	Large flake (FS54) manufactured on non-local banded rhyolite recovered from the surface of Square G on the quarry. Note the heavy edge-damage along the distal margin. Scale=1cm (Photograph: Paul Aurisch).	139
9.10	Water-rounded microgranite hammerstone (FS88) exhibiting impact-pitting recovered from the surface of Square J on the quarry. Several similar artefacts were noted eroding from adjacent midden deposits. Bustard Head, some 11km to the north is the nearest source of this raw material. Scale=1cm (Photograph: Paul Aurisch).	140
9.11	General view of completed excavation, Squares L–M. Facing southeast.	140
9.12	Close-up view of edge-ground hatchet (FS2747) (Photograph: Paul Aurisch).	140
9.13	Stratigraphic section, Ironbark Site Complex, Squares L–M.	142
9.14	Stratigraphic section, Ironbark Site Complex, Square N.	142
9.15	Stratigraphic section, Ironbark Site Complex, Squares O–P.	143
9.16	Stratigraphic section, Ironbark Site Complex, Squares Q–R.	144
9.17	Abundance of artefactual stone.	148
9.18	Abundance of artefactual stone.	149
9.19	Abundance of oyster (<i>S. glomerata</i>).	149
9.20	Abundance of oyster (<i>S. glomerata</i>).	149
9.21	Abundance of charcoal.	150
9.22	Abundance of charcoal.	150
9.23	Abundance of pumice.	150
9.24	Abundance of pumice.	151
9.25	Weight-depth distribution of individually provenanced artefacts greater than 30mm in maximum dimension, Square E (n=784).	153
10.1	The Eurimbula Creek catchment area showing the location of Eurimbula Creek 1 (EC1) and nearby excavated sites (ISC=Ironbark Site Complex; EC2=Eurimbula Creek 2; ES1=Eurimbula Site 1; TCSC=Tom’s Creek Site Complex). Dark grey shading indicates the general extent of mangrove, saltflats and claypans. Dotted shading indicates land above 200m. Solid dots indicate local population centres.	158
10.2	Site plan of Eurimbula Creek 1 area. Contours are in 0.5m intervals.	158
10.3	Excavations in progress at Squares A–D, showing cattle track on western (left) margin of the excavation. Facing north.	159
10.4	General view of completed excavation, Squares A–D, showing transect through erosion bank. Facing north.	159
10.5	General view of completed excavation, Squares A–D, showing transect through erosion bank. Facing east.	159
10.6	Stratigraphic section, Eurimbula Creek 1, Squares A–D.	161
10.7	Abundance of oyster (<i>S. glomerata</i>).	164
10.8	Abundance of oyster (<i>S. glomerata</i>).	164
10.9	Abundance of charcoal.	165
10.10	Abundance of charcoal.	165
10.11	Abundance of pumice.	165
10.12	Abundance of pumice.	166
11.1	Site plan of Eurimbula Creek 2 area. Contours are in 0.5m intervals. Only major trees in the immediate area of the excavation are shown.	170
11.2	General view of location of Square A, showing disturbance zone to the northeast (rear left) of the excavation. Facing east.	170
11.3	General view of completed excavation, Square A, showing a large root protruding from the east section. Facing north.	170
11.4	General view of completed excavation, showing disturbance zone to the east (right) of the excavation. Facing north.	170

11.5	Stratigraphic section, Eurimbula Creek 2, Square A.	172
11.6	Abundance of oyster (<i>S. glomerata</i>).	174
11.7	Abundance of charcoal.	175
11.8	Abundance of pumice.	175
11.9	Abundance of organic material.	175
12.1	Lower reaches of Round Hill Creek, showing the location of Transects A, B and C at Eurimbula Site 1. Dark grey shading indicates the general extent of mangrove vegetation. Stippling indicates the general location of intertidal flats. Dashed lines are sealed roads.	178
12.2	Location of test pits along Transects A, B and C at Eurimbula Site 1, showing topography in the immediate area of the transects.	182
12.3	General view of steep c.2m high erosion bank at the southern end of Eurimbula Site 1 fronting Round Hill Creek. Facing northwest.	182
12.4	Large stone artefact (FS1/2001) manufactured on rhyolitic tuff located mid-way down the erosion bank. Facing west.	182
12.5	Cleaned c.2.5m high section of erosion bank located along the southern third of the site, showing dark band of coffee rock mid-way down the profile. Facing west.	183
12.6	General view of completed excavation, Square E1, showing shell lens mid-way down the western profile. Facing west.	183
12.7	Close-up view of mud ark (<i>A. trapezia</i>) shell lens encountered during excavation, Square E1, XU10. Facing west.	183
12.8	General view of completed excavation at Squares A–D. Note large roots towards the top of the profile. Facing southwest.	183
12.9	Stratigraphic section, Eurimbula Site 1, Squares A–D.	186
12.10	Age-depth relationship of all radiocarbon determinations obtained at Eurimbula Site 1 (n=12). The linear regression shown only includes the six dates obtained on charcoal samples from Transect A, Squares A–D, E1–E2.	187
12.11	Abundance of artefactual stone.	190
12.12	Abundance of fish bone.	190
12.13	Abundance of oyster (<i>S. glomerata</i>).	191
12.14	Abundance of mud ark (<i>A. trapezia</i>).	191
12.15	Abundance of dominant shell taxa.	191
12.16	Abundance of organic material.	192
12.17	Abundance of charcoal.	192
12.18	Abundance of charcoal.	192
13.1	Aerial view of the Tom's Creek Site Complex area (after BPA Run 10B/57, 30 July 1996). The dashed box encloses the approximate area shown in Fig. 13.2. Based on data provided by the Department of Natural Resources and Mines, Queensland 2006, which gives no warranty in relation to the data (including accuracy, reliability, completeness or suitability) and accepts no liability (including without limitation, liability in negligence) for any loss, damage or costs (including consequential damage) relating to any use of the data.	203
13.2	Site plan of area of Squares A–S. Contours are in 0.5m intervals.	203
13.3	Schematic diagram of the layout of Squares A–D and E–Q, showing distribution of collected glass artefacts as solid dots (not to scale).	205
13.4	General view of area of glass scatter after removal of leaf litter. Flags indicate the position of glass artefacts. Facing northeast.	206
13.5	Glass artefact (FS188) cached in quinine tree. Facing north.	206
13.6	General view of completed excavation, Squares A–D. Facing northwest.	206

13.7	General view of shell material concentrated on erosion bank of Residual 1. Facing south.	207
13.8	General view of completed excavation, Squares R–S. Note position of shell material in the upper deposit. Facing north.	207
13.9	Core taken from below the limits of excavation, Square S. Note distinctive break (at c.75cm below ground surface) between coarse light yellow sands and dark organic muds.	207
13.10	Stratigraphic section, Tom’s Creek Site Complex, Squares A–D.	208
13.11	Stratigraphic section, Tom’s Creek Site Complex, Squares R–S.	209
13.12	Abundance of dominant shell taxa.	212
13.13	Abundance of dominant shell taxa.	215
13.14	Abundance of artefactual stone.	215
13.15	Abundance of artefactual stone.	215
13.16	Abundance of fish bone.	216
13.17	Abundance of fish bone.	216
13.18	Abundance of charcoal.	216
13.19	Abundance of charcoal.	217
14.1	Occupation spans of dated sites on the southern Curtis Coast, based on 1σ calibrated age-ranges. Note that a span of 100 years is estimated for the modern dates reported for WCM and EC2.	227
14.2	Estimated number of dated sites occupied on the southern Curtis Coast in each 500 year period, based on the mid-points of calibrated age-ranges.	228
14.3	Estimated number of new sites established on the southern Curtis Coast in each 500 year period. Note that the mid-point of the calibrated age-range of the oldest date available for each site is assumed to be the basal age.	228
14.4	Calibrated radiocarbon ages from the southern Curtis Coast (n=58) arranged in order of increasing age. Error bars show the 1σ calibrated age-range. Note the apparent gap between 1,050–1,250 years ago.	228
14.5	Number of sites on the southern Curtis Coast with central calibrated radiocarbon dates falling in each 500 year period, measured at 250 year intervals. For comparison, the same data are shown for all of southeast Queensland and the Moreton Bay region.	229
14.6	Summed probability plot of all calibrated radiocarbon ages (n=56) normalised to a maximum of one. Note that the two modern dates reported for WCM and EC2 are excluded.	229
14.7	Total weight of shell recovered from all excavated sites per 500 year interval. Note logarithmic scale.	234
14.8	Shellfish diversity calculated using the Shannon-Weaver Function (H’) per 500 year interval.	235
14.9	Total weight of fish bone recovered from all excavated sites per 500 year interval.	235
14.10	Total weight of charcoal recovered from all excavated sites per 500 year interval.	235
14.11	Total weight of stone artefacts recovered from all excavated sites per 500 year interval.	236
14.12	Abundance of local (line) versus non-local (black columns) stone artefact raw materials per 500 year interval.	236
14.13	Estimated site area for excavated sites.	239
14.14	Shell density as a unit of weight of excavated deposit for excavated sites. Only shell-bearing squares are included in calculations. Note logarithmic scale.	239
14.15	Shell density as a unit of area of excavated deposit.	239
14.16	Extrapolated total site shell content based on site area and shell density.	240
15.1	Southeast and central Queensland showing the location of major archaeological projects.	249

List of Tables

1.1	Average number of dates/site for various subsets of the radiocarbon dates available for archaeological sites in Queensland (Ulm and Reid 2000).	7
2.1	Creek/estuary characteristics in the study region (Olsen 1980a:17–25).	17
2.2	Radiocarbon dates from unexcavated sites in the study region (see Appendix 1 for full radiometric data for each determination). ABM=Agnes Beach Midden; MISS=Middle Island Sandblow Site; RHCM=Round Hill Creek Mound; WCM=Worthington Creek Midden. Dates on shell were calibrated using a ΔR value of $+10\pm 7$, except Wk-10090 where $\Delta R = -155\pm 55$ (see Chapter 4 for further details).	30
2.3	Middle Island Sandblow Site, Squares A–E: summary excavation data and dominant materials.	33
3.1	Summary of excavated sites (arranged north to south).	38
4.1	Original 1970s series of radiocarbon dates obtained on live-collected marine shell samples from Australian waters presented by Gillespie (1977; Gillespie and Temple 1977). $\delta^{13}\text{C}$ is an assumed value of 0 ± 2 (Gillespie and Polach 1979:410). Historical ages of shell samples were converted to equivalent global marine model ages using data from Stuiver et al. (1998a). ΔR was calculated by deducting the equivalent marine model age of the historical age of the shell sample from the ^{14}C age of the shell sample (after Stuiver et al. 1986:1020). $\Delta R\sigma = \sqrt{(\sigma_{\text{historical age}}^2 + \sigma_{\text{marine model age}}^2 + \sigma_{^{14}\text{C age}}^2)}$ (Gillespie 1982). The uncertainty in the marine model age includes estimated error in the calibration dataset (derived from Stuiver et al. 1998a). Error-weighted means are calculated using formulae in Ward and Wilson (1978). Samples: Mo= <i>Mactra obesa</i> ; Pb= <i>Pinna bicolor</i> ; Pm= <i>Pinctada margaritifera</i> ; Pl= <i>Proxichione laqueata</i> ; Dd= <i>Donax deltoides</i> ; Kr= <i>Katelysia rhytiphora</i> .	52
4.2	Radiocarbon dates obtained on live-collected marine shell samples of known historical age from the Gulf of Carpentaria (Rhodes et al. 1980). Samples: A= <i>Anadara</i> sp.; Tt= <i>Telescopium telescopium</i> . See caption for Table 4.1 for details of calculations.	53
4.3	ΔR values for Abraham Reef and Heron Island coral cores (after Reimer and Reimer 2000). Samples: <i>Porites australiensis</i> . See caption for Table 4.1 for details of calculations.	54
4.4	Post-AD 1950 live-collected shell (Gillespie and Polach 1979:Table 5; Rhodes et al. 1980:Table 1). Samples: Mep= <i>Mytilus edulis planulatus</i> ; Pe= <i>Pyrazus ebeninus</i> ; V= <i>Volachlamys</i> sp.; Ss= <i>Saccostrea succulata</i> .	55
4.5	Radiocarbon ages obtained on 'pre-bomb' live-collected marine shell samples from central Queensland. Samples: Dd= <i>Donax deltoides</i> ; At= <i>Anadara trapezia</i> ; Vs= <i>Volachlamys singaporina</i> . See caption for Table 4.1 for details of calculations.	55
4.6	Shell/charcoal paired samples from the southern Curtis Coast. ^{14}C ages obtained on charcoal samples were reduced by 41 ± 14 years to correct for ^{14}C variation between and northern southern hemispheres (McCormac et al. 2002). An estimate of the atmospheric calibration curve error, derived from an average of estimated error in the 1σ span of the age, was also included. Therefore, atmospheric age $\sigma = \sqrt{(\sigma_{^{14}\text{C age}}^2 + \sigma_{\text{southern hemisphere offset}}^2 + \text{average of calibration curve error}^2)}$ (Gillespie 1982). Note that the incorporation of southern hemisphere offset error in this formula assumes that each atmospheric conventional radiocarbon age derives from an independent secondary reservoir (see Jones and Nicholls 2001 for discussion). The 1σ range of the ^{14}C value was converted to the equivalent global marine model 1σ range using atmospheric ages interpolated from INTCAL98 to the same calendar year as MARINE98 (Stuiver et al. 1998a). ΔR was	59

calculated by deducting the mid-point of the equivalent marine model age of the charcoal determination from the ^{14}C age of the paired marine shell sample. $\Delta R\sigma = \sqrt{(\sigma_{\text{marine model age}}^2 + \sigma_{\text{marine shell age}}^2)}$ (Gillespie 1982). This method is illustrated for pair NZA-12117/Wk-8326 in Fig. 4.3.

4.7	Calibrated radiocarbon ages from the Seven Mile Creek Mound, using various ΔR values.	63
5.1	Attributes of articulated <i>A. trapezia</i> specimens recovered from the Seven Mile Creek Mound in rank order of correlation coefficient. Note that specimens were only included if there was no damage inhibiting accurate measurement of each attribute on either valve in a pair.	68
5.2	Attributes of live-collected <i>A. trapezia</i> specimens in rank order of correlation coefficient.	68
5.3	Identified <i>A. trapezia</i> conjoin sets, Seven Mile Creek Mound, Square A.	73
5.4	Additional <i>A. trapezia</i> conjoin sets identified during the blind test, Seven Mile Creek Mound, Square A.	74
5.5	Summary of maximum distance separating all conjoined <i>A. trapezia</i> valve-pairs, Seven Mile Creek Mound, Square A.	74
6.1	Stratigraphic Unit descriptions, Seven Mile Creek Mound, Squares A–D.	83
6.2	Radiocarbon dates from the Seven Mile Creek Mound, Square A (see Appendix 1 for full radiometric data for each determination).	84
6.3	Presence/absence of shellfish identified in the Seven Mile Creek Mound, Square A.	88
6.4	Metrical data for intact and broken (with umbo) <i>A. trapezia</i> valves from the Seven Mile Creek Mound, Square A.	91
6.5	Fish bone abundance, Seven Mile Creek Mound, Square A.	93
6.6	Fish bone taxonomic representation, Seven Mile Creek Mound, Square A.	93
6.7	Stone artefacts from the Seven Mile Creek Mound, Square A.	94
7.1	Stratigraphic Unit descriptions, Mort Creek Site Complex, Squares A–D.	103
7.2	Radiocarbon dates from the Mort Creek Site Complex (see Appendix 1 for full radiometric data for each determination).	105
7.3	Identified <i>A. trapezia</i> conjoin sets, Mort Creek Site Complex, Square C.	105
7.4	Presence/absence of shellfish identified in the Mort Creek Site Complex, Square C.	107
7.5	Metrical data for intact and broken (with umbo) <i>A. trapezia</i> valves from the Mort Creek Site Complex, Square C.	110
7.6	Fish bone abundance, Mort Creek Site Complex, Square C.	111
7.7	Fish bone taxonomic representation, Mort Creek Site Complex, Square C.	111
7.8	Stone artefacts from the Mort Creek Site Complex, Square C.	111
8.1	Pancake Creek Site Complex, Squares A–H: summary excavation data and dominant materials.	120
8.2	Stratigraphic Unit descriptions, Pancake Creek Site Complex, Squares A–B.	120
8.3	Stratigraphic Unit descriptions, Pancake Creek Site Complex, Squares C–F.	121
8.4	Stratigraphic Unit descriptions, Pancake Creek Site Complex, Squares G–H.	122
8.5	Radiocarbon dates from the Pancake Creek Site Complex (see Appendix 1 for full radiometric data for each determination).	123
8.6	Presence/absence of shellfish identified in the Pancake Creek Site Complex, Squares A–H.	125
8.7	Stone artefacts from the Pancake Creek Site Complex, Squares A–H.	128
9.1	Ironbark Site Complex, Squares E–R: summary excavation data and dominant materials.	141
9.2	Stratigraphic Unit descriptions, Ironbark Site Complex, Square E.	141

9.3	Stratigraphic Unit descriptions, Ironbark Site Complex, Squares L–M.	141
9.4	Stratigraphic Unit descriptions, Ironbark Site Complex, Square N.	142
9.5	Stratigraphic Unit descriptions, Ironbark Site Complex, Squares O–P.	143
9.6	Stratigraphic Unit descriptions, Ironbark Site Complex, Squares Q–R.	144
9.7	Radiocarbon dates from the Ironbark Site Complex (see Appendix 1 for full radiometric data for each determination). * = assumed value only.	145
9.8	Presence/absence of shellfish identified in the Ironbark Site Complex, Squares L–R.	148
9.9	Fish bone abundance, Ironbark Site Complex, Squares N, O and P.	151
9.10	Stone artefacts from the Ironbark Site Complex, Squares E, L–R.	153
10.1	Eurimbula Creek 1, Squares A–D: summary excavation data and dominant materials.	161
10.2	Stratigraphic Unit descriptions, Eurimbula Creek 1, Squares A–D.	161
10.3	Radiocarbon dates from Eurimbula Creek 1 (see Appendix 1 for full radiometric data).	162
10.4	Presence/absence of shellfish identified in Eurimbula Creek 1, Squares A–D.	163
10.5	Fish bone abundance, Eurimbula Creek 1, Squares A–D.	164
11.1	Eurimbula Creek 2, Square A: summary excavation data and dominant materials.	172
11.2	Stratigraphic Unit descriptions, Eurimbula Creek 2, Square A.	172
11.3	Radiocarbon dates from Eurimbula Creek 2 (see Appendix 1 for full radiometric data).	173
11.4	Presence/absence of shellfish identified in Eurimbula Creek 2, Square A.	174
12.1	Fish bone abundance, Eurimbula Site 1, Squares E1–E2.	183
12.2	Eurimbula Site 1, Squares A–D: summary excavation data and dominant materials. Data from Squares E1–E9 are included for comparison (after Ulm et al. 1999a:Appendix A–I).	185
12.3	Stratigraphic Unit descriptions, Eurimbula Site 1, Squares A–D.	185
12.4	Radiocarbon dates from Eurimbula Site 1 (see Appendix 1 for full radiometric data for each determination). E* This date was undertaken on a sample of shell from a dense surface scatter adjacent to Square E7.	187
12.5	Identified <i>A. trapezia</i> conjoin sets, Eurimbula Site 1, Squares A–D.	189
12.6	Presence/absence of shellfish identified in Eurimbula Site 1, Squares A–D.	193
12.7	Metrical data for intact and broken (with umbo) <i>A. trapezia</i> valves from Eurimbula Site 1, Squares A–D. Note that excavation units for each square have been collapsed for purposes of analysis. Excavation unit depth and size is approximately equivalent across squares for broad comparison.	194
12.8	Fish bone abundance, Eurimbula Site 1, Squares A–D.	195
12.9	Stone artefacts from Eurimbula Site 1, Squares A–D.	197
13.1	Tom’s Creek Site Complex, Squares A–D, R–S: summary excavation data and dominant materials.	208
13.2	Stratigraphic Unit descriptions, Tom’s Creek Site Complex, Squares A–D.	208
13.3	Stratigraphic Unit descriptions, Tom’s Creek Site Complex, Squares R–S.	209
13.4	Radiocarbon dates from the Tom’s Creek Site Complex (see Appendix 1 for full radiometric data for each determination).	210
13.5	Identified <i>A. trapezia</i> conjoin sets, Tom’s Creek Site Complex, Squares A–D.	211
13.6	Identified <i>A. trapezia</i> conjoin sets, Tom’s Creek Site Complex, Squares R–S.	211
13.7	Presence/absence of shellfish identified in the Tom’s Creek Site Complex, Squares A–S.	213
13.8	Metrical data for intact and broken (with umbo) <i>A. trapezia</i> valves from the Tom’s Creek Site Complex, Squares A–D. Note that excavation units for each square have been collapsed for purposes of analysis. Excavation unit depth and size is approximately equivalent across squares for broad comparison.	217

13.9	Metrical data for intact and broken (with umbo) <i>A. trapezia</i> valves from the Tom's Creek Site Complex, Squares R–S. Note that excavation units for each square have been collapsed for purposes of analysis. Excavation unit depth and size is approximately equivalent across squares for broad comparison.	218
13.10	Fish bone abundance, Tom's Creek Site Complex, Squares A–D.	219
13.11	Fish bone abundance, Tom's Creek Site Complex, Squares R–S.	220
13.12	Stone artefacts from the Tom's Creek Site Complex, Squares A–D, R–S.	221
13.13	Glass artefacts from the Tom's Creek Site Complex. * indicates glass artefacts recovered beyond the mapping grid (see Fig. 13.3).	222
14.1	Radiocarbon dates from the southern Curtis Coast excluded from chronological analyses.	227
14.2	Summed probability distribution of all calibrated radiocarbon dates available from cultural contexts on the southern Curtis Coast (n=56). Excludes the two modern determinations (Wk-7681; Wk-7689). 0* indicates a modern age.	231
14.3	Radiocarbon ages from the southern Curtis Coast dating to the 1,500–2,000 cal BP interval, calibrated using various ΔR values.	232
14.4	Top 10 shellfish taxa from all excavated deposits ranked by weight and minimum number of individuals. Note that the four taxa of terrestrial gastropod have been excluded.	233
14.5	Shell density characteristics of excavated sites arranged in descending order of total estimated shell content.	238